

- ✓ Page 19, line 13, please replace "acdi" with --acid--.
- ✓ Page 21, line 29, please replace "skilied" with --skilled--;  
✓ line 30, please replace "enyzmatic" with --enzymatic--.
- ✓ Page 25, line 32, please replace "effecting" with --affecting--.
- ✓ Page 28, line 13, please replace "spleen.y" with --spleen.--.
- ✓ Page 31, line 6, after "48" please insert --and--;  
✓ line 22, please replace "ng/mg" with --ng/mg)--;  
✓ line 26, please replace "ng/ml" with --ng/ml)--.
- Page 32, line 11, please replace "meonatal" with --neonatal--;  
✓ line 26, please replace "Shingomyelin(SM):EPC:CHOL" with  
--Sphingomyelin(SM):EPC:CHOL--;
- ✓ Page 33, line 1, please replace "of<sup>32</sup>P" with --of <sup>32</sup>P--;  
line 2, between "i.v." and "made," please insert --was--.

IN THE CLAIMS:

✓ Please cancel claims 1-61.

Please add new claims 62-83:

---

1                    ~~--62.~~ A method of treating a neoplasia in a mammal, said method comprising  
2     administering to said mammal a pharmaceutical composition comprising a polyethylene glycol  
3     (PEG)-ceramide conjugate, a lipid, and a nucleic acid catalyst, whereby a therapeutic result is  
4     attained.

1                    63. The method of claim 62, wherein the administration of said composition  
2     results in a decrease in the expression of an RNA associated with said neoplasia in said  
3     mammal.

1                    64. The method of claim 62, wherein said neoplasia comprises a tumor.

1                    65. The method of claim 62, wherein the administration is a systemic  
2     administration.

1                   66. The method of claim 62, wherein said composition further comprises  
2 phosphatidyl choline.

1                   67. The method of claim 66, wherein said phosphatidyl choline is egg yolk  
2 phosphatidyl choline.

1                   68. The method of claim 62, wherein said composition further comprises  
2 cholesterol.

1                   69. The method of claim 62, wherein said nucleic acid catalyst has an  
2 endonuclease activity.

1                   70. The method of claim 62, wherein said nucleic acid catalyst comprises a  
2 ribonucleotide or a deoxyribonucleotide.

1                   71. The method of claim 62, wherein said nucleic acid catalyst comprises a  
2 hammerhead motif.

1                   72. The method of claim 62, wherein said lipid is a cationic lipid.

1                   73. The method of claim 72, wherein said cationic lipid is DODAC or DOTAP.

1                   74. The method of claim 62, wherein said PEG-ceramide conjugate comprises a  
2 fatty acid group having eight carbon atoms.

1                   75. The method of claim 62, wherein said PEG-ceramide conjugate comprises a  
2 fatty acid group having fourteen carbon atoms.

1                   76. The method of claim 62, wherein said PEG-ceramide conjugate comprises a  
2 fatty acid group having twenty carbon atoms.

1                   77. The method of claim 62, wherein said nucleic acid catalyst specifically  
2 cleaves vascular endothelial growth factor receptor (VEGF-R) RNA.

a  
cont.